Updated on Apr-09



## **LavaDigest Interfering Compounds**

## Acceptable maximum levels of chemicals for LavaDigest kit.

The acceptable upper limit of interfering compounds commonly used in preparation of protein samples for digestions is provided in the table below.

| Chemicals                       | Manufacturer  | Recommended limit of chemicals found in 100 µL of protein samples in the LavaDigest assay |
|---------------------------------|---------------|---|
| SDS                             | BDH           | 0.02%   |
| triton X 100                    | UT Packard    | 0.001% v/v  |
| tween 20                        | Sigma         | 0.002% v/v  |
| Guanidine-HCI                   | BDH           | 500 μΜ  |
| CHAPS                           | BioRad        | 0.02%   |
| NaOH                            | BDH           | 10 mM   |
| dithiothreitol                  | BioRad        | 4 mM  |
| iodoacetamide                   | Sigma-Aldrich | 3 mM  |
| Urea                            | Sigma-Aldrich | 1 M   |
| ca <mark>lciu</mark> m chloride | BDH           | 5 mM  |
| tris-HCI                        | Sigma-Aldrich | 2.5 mM  |

Acceptable maximum was defined as difference in the constant decay rate (k2) driven from trypsin-digested BSA sample less than 20% coefficient of variance between control and the tests containing interfering chemicals.

